

Mountain pass solutions for a mean field equation from two-dimensional turbulence

Tonia Ricciardi*

Dipartimento di Matematica e Applicazioni “R. Caccioppoli”

Università di Napoli Federico II

Via Cintia, 80126 Napoli, Italy

fax: +39 081 675665

tonia.ricciardi@unina.it

December 5, 2006

Abstract

Using Struwe’s “monotonicity trick” and the recent blow-up analysis of Ohtsuka and Suzuki, we prove the existence of mountain pass solutions to a mean field equation arising in two-dimensional turbulence.

KEY WORDS: mean field equation, mountain pass solution

MSC 2000 SUBJECT CLASSIFICATION: 35J60

1 Introduction and main result

We consider the following problem:

$$(1) \quad \begin{aligned} -\Delta_g u &= \lambda_1 \left(\frac{e^u}{\int_M e^u dv_g} - \frac{1}{|M|} \right) - \lambda_2 \left(\frac{e^{-u}}{\int_M e^{-u} dv_g} - \frac{1}{|M|} \right) \\ \int_M u dv_g &= 0, \end{aligned}$$

where (M, g) is a compact, orientable, Riemannian 2-manifold without boundary, Δ_g denotes the Laplace-Beltrami operator, $|M|$ denotes the volume of M and where λ_1, λ_2 are positive constants. Equation (1) arises in the context of the statistical mechanics description of two-dimensional turbulence, see Joyce and Montgomery [4] and Pointin and Lundgren [7]. Problem (1) has a variational structure. Indeed, solutions to (1) correspond to critical points for the functional

$$\begin{aligned} I_{\lambda_1, \lambda_2}(u) &= \\ &= \frac{1}{2} \int_M |\nabla_g u|^2 dv_g - \lambda_1 \ln \left(\frac{1}{|M|} \int_M e^u dv_g \right) - \lambda_2 \ln \left(\frac{1}{|M|} \int_M e^{-u} dv_g \right) \end{aligned}$$

*Supported in part by the MIUR National Project *Variational Methods and Nonlinear Differential Equations*.

defined on the Sobolev space

$$E = \{u \in H^1(M) : \int_M u \, dv_g = 0\},$$

equipped with the norm $\|u\|^2 = \int_M |\nabla_g u|^2 \, dv_g$. A detailed analysis of the functional I_{λ_1, λ_2} has been recently carried out by Ohtsuka and Suzuki [6], together with a blow-up analysis for problem (1). In particular, it is shown in [6] that I_{λ_1, λ_2} is bounded below if and only if $0 \leq \lambda_1, \lambda_2 \leq 8\pi$, thus improving the classical sharp Moser-Trudinger inequality. The corresponding problem under Dirichlet boundary conditions has been recently considered in Sawada, Suzuki and Takahashi [8].

We note that (1) always admits the trivial solution $u \equiv 0$. Our aim in this note is to prove the existence of nontrivial solutions to problem (1) for suitable values of λ_1, λ_2 . We denote by $\mu_1(M)$ the first nonzero eigenvalue of Δ_g , namely,

$$\mu_1(M) = \inf_{u \in E \setminus \{0\}} \frac{\int_M |\nabla_g u|^2 \, dv_g}{\int_M u^2 \, dv_g}.$$

We assume that $\mu_1(M)$ satisfies the condition

$$(2) \quad 8\pi < \mu_1(M)|M| < 16\pi.$$

Condition (2) is satisfied, e.g., on the flat torus $\mathbb{R}^2/\mathbb{Z}^2$, since $\mu_1(\mathbb{R}^2/\mathbb{Z}^2) = 4\pi^2$. We define

$$\Lambda = \{(\lambda_1, \lambda_2) \in \mathbb{R}_0^+ \times \mathbb{R}_0^+ : \lambda_1 + \lambda_2 < \mu_1(M)|M| \text{ and } \max\{\lambda_1, \lambda_2\} > 8\pi\}.$$

Note that condition (2) ensures that $\Lambda \neq \emptyset$. In fact, Λ is the union of the triangle with vertices $(8\pi, 0)$, $(\mu_1(M)|M|, 0)$, $(8\pi, \mu_1(M)|M| - 8\pi)$, and of its reflection with respect to the straight line $\lambda_1 = \lambda_2$. We prove the following result:

Theorem 1. *Suppose (M, g) satisfies condition (2). Then problem (1) admits a nonzero solution for every $(\lambda_1, \lambda_2) \in \Lambda$.*

We note that when $\lambda_2 = 0$, Theorem 1 reduces to Theorem 2.1 obtained by Struwe and Tarantello in [10]. In fact, the proof of Theorem 1 employs the variational approach introduced in [10] and further developed by Jeanjean in [3], together with a fine blow-up analysis for solutions to (1) by Ohtsuka and Suzuki [6]. More precisely, in Section 2 we show that for every $(\lambda_1, \lambda_2) \in \Lambda$ the functional I_{λ_1, λ_2} has a mountain pass structure. However, since $\max\{\lambda_1, \lambda_2\} > 8\pi$ we have a lack of compactness, and general Palais-Smale sequences may “blow-up” in the sense of Brezis and Merle [1]. Therefore, in Section 3 we employ Struwe’s “monotonicity trick” [9] in order to find a bounded Palais-Smale sequence for almost every $(\lambda_1, \lambda_2) \in \Lambda$, and consequently a solution to (1). Finally, we exploit the blow-up analysis of Ohtsuka and Suzuki [6] to obtain solutions for every $(\lambda_1, \lambda_2) \in \Lambda$. Related results in the context of $SU(3)$ Toda systems may be found in Chae, Ohtsuka and Suzuki [2] and in Lucia and Nolasco [5].

2 Some properties of I_{λ_1, λ_2}

In this section we prove some properties of I_{λ_1, λ_2} which will be useful in what follows. In particular, we prove the following result:

Proposition 1. *The functional I_{λ_1, λ_2} has a mountain pass structure for every $(\lambda_1, \lambda_2) \in \Lambda$.*

Before proving Proposition 1 we introduce some notation and we establish some preliminary results. For every $u \in E$ we set

$$(3) \quad G(u) = \ln \left(\frac{1}{|M|} \int_M e^u dv_g \right).$$

Then, we may write

$$I_{\lambda_1, \lambda_2}(u) = \frac{1}{2} \int_M |\nabla_g u|^2 dv_g - \lambda_1 G(u) - \lambda_2 G(-u).$$

For every $\phi, \psi \in E$ we have

$$\begin{aligned} G'(u)\phi &= \frac{\int_M e^u \phi dv_g}{\int_M e^u dv_g} \\ \langle G''(u)\phi, \psi \rangle &= \frac{\int_M e^u \phi \psi dv_g \int_M e^u dv_g - \int_M e^u \phi dv_g \int_M e^u \psi dv_g}{\left(\int_M e^u dv_g \right)^2}. \end{aligned}$$

In particular, recalling that $\int_M \phi dv_g = 0$, we obtain

$$(4) \quad G'(0) = 0, \quad \langle G''(0)\phi, \phi \rangle = \frac{\int_M \phi^2 dv_g}{|M|}.$$

Furthermore, by the Cauchy-Schwarz inequality, we have

$$\left(\int_M e^u \phi dv_g \right)^2 \leq \int_M e^u \phi^2 dv_g \int_M e^u dv_g$$

for every $u, \phi \in E$, and therefore

$$(5) \quad \langle G''(u)\phi, \phi \rangle \geq 0 \quad \forall u, \phi \in E.$$

We shall also use the following property of G :

Lemma 1. *There exists a constant $C_M > 0$ depending on M only, such that*

$$\|G'(u)\| \leq C_M \exp \left\{ \frac{1}{8\pi} \|u\|^2 \right\}$$

for all $u \in E$.

Proof. By Jensen's inequality we have $\int_M e^u dv_g \geq |M|$. Therefore, for every $\phi \in E$ we have:

$$|G'(u)\phi| \leq \frac{1}{|M|} \int_M e^u \phi dv_g \leq \frac{1}{|M|} \left(\int_M e^{2u} dv_g \right)^{1/2} \left(\int_M \phi^2 dv_g \right)^{1/2}.$$

In view of the Moser-Trudinger inequality:

$$(6) \quad \int e^u dv_g \leq C_0 \exp \left\{ \frac{1}{16\pi} \|u\|^2 \right\} \quad \forall u \in E,$$

where $C_0 > 0$ depends on M only, and of the Poicaré inequality

$$(7) \quad \int_M \phi^2 dv_g \leq \frac{1}{\mu_1(M)} \int_M |\nabla_g \phi|^2 dv_g \quad \forall \phi \in E,$$

we derive

$$|G'(u)\phi| \leq \frac{[C_0 \exp\{\frac{1}{4\pi}\|u\|^2\}]^{1/2} \|\phi\|}{\mu_1(M)^{1/2}|M|}$$

and the asserted estimate follows with $C_M = (C_0/\mu_1(M))^{1/2} |M|^{-1}$. $\square$

Lemma 2. *For every $\lambda_1, \lambda_2 \geq 0$ such that $\lambda_1 + \lambda_2 < \mu_1(M)|M|$ the function $u \equiv 0$ is a local minimum for I_{λ_1, λ_2} .*

Proof. We have:

$$\begin{aligned} I_{\lambda_1, \lambda_2}(0) &= 0, & I'_{\lambda_1, \lambda_2}(0) &= 0, \\ \langle I''_{\lambda_1, \lambda_2}(0)\phi, \phi \rangle &= \int_M |\nabla_g \phi|^2 dv_g - \frac{\lambda_1 + \lambda_2}{|M|} \int_M \phi^2 dv_g. \end{aligned}$$

In view of the Poicaré inequality (7) we derive

$$\langle I''_{\lambda_1, \lambda_2}(0)\phi, \phi \rangle \geq \left(1 - \frac{\lambda_1 + \lambda_2}{\mu_1(M)|M|}\right) \|\phi\|^2.$$

Now the claim follows by Taylor expansion of I_{λ_1, λ_2} at $u \equiv 0$. $\square$

Let $p_0 \in M$ and let $r_0 > 0$ be a constant smaller than the injectivity radius of M at p_0 . Let $\mathcal{B}_{r_0} = \{p \in M : d_g(p, p_0) < r_0\}$ denote the geodesic ball of radius r_0 centered at p_0 . For every $\varepsilon > 0$ let v_ε be the function defined by

$$v_\varepsilon(p) = \begin{cases} \ln \frac{\varepsilon^2}{(\varepsilon^2 + d_g(p, p_0)^2)^2} & \text{in } \mathcal{B}_{r_0} \\ \ln \frac{\varepsilon^2}{(\varepsilon^2 + r_0^2)^2} & \text{in } M \setminus \mathcal{B}_{r_0} \end{cases}$$

and let $u_\varepsilon \in E$ be correspondingly defined by

$$u_\varepsilon = v_\varepsilon - \frac{1}{|M|} \int_M v_\varepsilon dv_g.$$

We have the following asymptotic expansions.

Lemma 3. *There exists $r_0 > 0$ such that the following asymptotic expansions hold, with respect to $\varepsilon \rightarrow 0$:*

$$(8) \quad \int_M |\nabla_g u_\varepsilon|^2 dv_g = 32\pi \ln \frac{1}{\varepsilon} + O(1)$$

$$(9) \quad \ln \int_M e^{u_\varepsilon} dv_g = 2 \ln \frac{1}{\varepsilon} + O(1)$$

$$(10) \quad \ln \int_M e^{-u_\varepsilon} dv_g = O(1).$$

Proof. In geodesic coordinates centered at p we have $g_{ij}(x) = \delta_{ij} + O(|x|^2)$ for every $x \in B_{r_0} = \{x \in \mathbb{R}^2 : |x| < r_0\}$. Consequently, $dv_g = (1 + O(|x|^2)) dx$ and $|\nabla_g u|^2 dv_g = |\nabla u|^2 (1 + O(|x|^2)) dx$, where ∇ denotes differentiation with respect to the Euclidean metric. Moreover, identifying $v_\varepsilon, u_\varepsilon$ with their pullbacks to B_{r_0} under the geodesic coordinate system, we have in B_{r_0} :

$$v_\varepsilon(x) = \ln \frac{\varepsilon^2}{(\varepsilon^2 + |x|^2)^2},$$

$$\nabla u_\varepsilon(x) = \nabla v_\varepsilon(x) = -\frac{4x}{\varepsilon^2 + |x|^2}.$$

Proof of (8). We have

$$\begin{aligned} \int_{B_{r_0}} |\nabla u_\varepsilon|^2 dx &= 32\pi \int_0^{r_0} \frac{r^3}{(\varepsilon^2 + r^2)^2} dr \\ &= 32\pi \left(\int_0^{r_0} \frac{r dr}{\varepsilon^2 + r^2} - \varepsilon^2 \int_0^{r_0} \frac{r dr}{(\varepsilon^2 + r^2)^2} \right) \\ &= 16\pi \left(\ln(\varepsilon^2 + r^2) \Big|_0^{r_0} + \frac{\varepsilon^2}{\varepsilon^2 + r^2} \Big|_0^{r_0} \right) = 16\pi \ln \varepsilon^{-2} + O(1). \end{aligned}$$

Moreover,

$$\int_{B_{r_0}} |\nabla u_\varepsilon|^2 |x|^2 dx \leq 16 \int_{B_{r_0}} dx = O(1).$$

Since $\nabla_g u_\varepsilon \equiv 0$ in $M \setminus \mathcal{B}_{r_0}$, we conclude that

$$\int_M |\nabla_g u_\varepsilon|^2 dv_g = \int_{B_{r_0}} |\nabla u_\varepsilon|^2 (1 + O(|x|^2)) dx = 16\pi \ln \varepsilon^{-2} + O(1)$$

and (8) is established.

Proof of (9). We have:

$$\begin{aligned} \int_{B_{r_0}} e^{v_\varepsilon} dx &= \int_{B_{r_0}} \frac{\varepsilon^2}{(\varepsilon^2 + |x|^2)^2} dx = \int_{B_{r_0/\varepsilon}} \frac{dy}{(1 + |y|^2)^2} = \int_{\mathbb{R}^2} \frac{dy}{(1 + |y|^2)^2} + o(1) \\ \int_{B_{r_0}} e^{v_\varepsilon} |x|^2 dx &= \int_{B_{r_0}} \frac{\varepsilon^2 |x|^2}{(\varepsilon^2 + |x|^2)^2} dx = \varepsilon^2 \int_{B_{r_0/\varepsilon}} \frac{|y|^2 dy}{(1 + |y|^2)^2} = o(1), \\ \int_{M \setminus \mathcal{B}_{r_0}} e^{v_\varepsilon} dv_g &= \frac{\varepsilon^2}{(\varepsilon^2 + r_0^2)} |M \setminus \mathcal{B}_{r_0}| = o(1). \end{aligned}$$

Therefore, we obtain

$$\int_M e^{v_\varepsilon} dv_g = \int_{B_{r_0}} e^{v_\varepsilon} (1 + O(|x|^2)) dx + \int_{M \setminus \mathcal{B}_{r_0}} e^{v_\varepsilon} dv_g = \int_{\mathbb{R}^2} \frac{dy}{(1 + |y|^2)^2} + o(1)$$

and consequently

$$(11) \quad \ln \int_M e^{v_\varepsilon} dv_g = \ln \left(\int_{\mathbb{R}^2} \frac{dy}{(1 + |y|^2)^2} + o(1) \right) = O(1).$$

Moreover, we have

$$\begin{aligned}\int_M v_\varepsilon dv_g &= |M| \ln \varepsilon^2 - 2 \int_{B_{r_0}} \ln(\varepsilon^2 + |x|^2) dv_g - 2|M \setminus \mathcal{B}_{r_0}| \ln(\varepsilon^2 + r_0^2) \\ &= |M| \ln \varepsilon^2 + O(1)\end{aligned}$$

and therefore the mean value of v_ε satisfies

$$(12) \quad \frac{1}{|M|} \int_M v_\varepsilon dv_g = \ln \varepsilon^2 + O(1).$$

Consequently, in view of (11) and (12),

$$\ln \int_M e^{u_\varepsilon} dv_g = \ln \int_M e^{v_\varepsilon} dv_g - \frac{1}{|M|} \int_M v_\varepsilon dv_g = \ln \varepsilon^{-2} + O(1).$$

Hence, (9) is established.

Proof of (10). Using the elementary formula $\int_{B_{r_0}} |x|^p dx = 2\pi(p+2)^{-1}r_0^{p+2}$, we compute:

$$\begin{aligned}\int_{B_{r_0}} e^{-v_\varepsilon} dx &= \int_{B_{r_0}} \frac{(\varepsilon^2 + |x|^2)^2}{\varepsilon^2} dx \\ &= \varepsilon^{-2} \int_{B_{r_0}} |x|^4 dx + 2 \int_{B_{r_0}} |x|^2 dx + \varepsilon^2 \int_{B_{r_0}} dx \\ &= \frac{\pi r_0^6}{3} \varepsilon^{-2} + \pi r_0^4 + \varepsilon^2 \pi r_0^2, \\ \int_{B_{r_0}} e^{-v_\varepsilon} |x|^2 dx &= \varepsilon^{-2} \int_{B_{r_0}} |x|^6 dx + 2 \int_{B_{r_0}} |x|^4 dx + \varepsilon^2 \int_{B_{r_0}} |x|^2 dx \\ &= \frac{\pi r_0^8}{4} \varepsilon^{-2} + \frac{2\pi r_0^6}{3} + \varepsilon^2 \frac{\pi r_0^4}{2}.\end{aligned}$$

Therefore,

$$\begin{aligned}\int_{B_{r_0}} e^{-v_\varepsilon} dv_g &= \int_{B_{r_0}} \frac{(\varepsilon^2 + |x|^2)^2}{\varepsilon^2} (1 + O(|x|^2)) dx \\ &= \frac{\pi r_0^6}{3} (1 + O(r_0)^2) \varepsilon^{-2} + \pi r_0^4 (1 + O(r_0)^2) + \varepsilon^2 \pi r_0^2 (1 + O(r_0)^2)\end{aligned}$$

and by choosing $r_0 > 0$ sufficiently small, we obtain

$$\int_{B_{r_0}} e^{-v_\varepsilon} dv_g = \beta_0 \varepsilon^{-2} + \beta_1 + o(1)$$

for some $\beta_0, \beta_1 > 0$. On the other hand, in $M \setminus \mathcal{B}_{r_0}$ we have

$$\begin{aligned}\int_{M \setminus \mathcal{B}_{r_0}} e^{-v_\varepsilon} dv_g &= \frac{(\varepsilon^2 + r_0^2)^2}{\varepsilon^2} |M \setminus \mathcal{B}_{r_0}| \\ &= |M \setminus \mathcal{B}_{r_0}| \varepsilon^{-2} + 2r_0^2 |M \setminus \mathcal{B}_{r_0}| + o(1).\end{aligned}$$

It follows that

$$\int_M e^{-v_\varepsilon} dv_g = (\beta_0 + |M \setminus \mathcal{B}_{r_0}|) \varepsilon^{-2} + \beta_1 + 2r_0^2 |M \setminus \mathcal{B}_{r_0}| + o(1)$$

and consequently,

$$\begin{aligned}\ln \int_M e^{-v_\varepsilon} dv_g &= \ln [(\beta_0 + |M \setminus \mathcal{B}_{r_0}|)\varepsilon^{-2} + \beta_1 + 2r_0^2|M \setminus \mathcal{B}_{r_0}| + o(1)] \\ &= \ln \varepsilon^{-2} + \ln [\beta_0 + |M \setminus \mathcal{B}_{r_0}| + o(1)] = \ln \varepsilon^{-2} + O(1).\end{aligned}$$

In view of the above and (12), we derive

$$\begin{aligned}\ln \int_M e^{-u_\varepsilon} dv_g &= \ln \int_M e^{-v_\varepsilon} dv_g + \frac{1}{|M|} \int_M v_\varepsilon dv_g \\ &= \ln \varepsilon^{-2} + \ln \varepsilon^2 + O(1) = O(1).\end{aligned}$$

This establishes (10). $\square$

Proof of Proposition 1. Let $(\lambda_1, \lambda_2) \in \Lambda$. In view of Lemma 2, $u \equiv 0$ is a strict nondegenerate local minimum for I_{λ_1, λ_2} . Moreover, in view of the expansions (8), (9) and (10), we have

$$I_{\lambda_1, \lambda_2}(u_\varepsilon) = 2(8\pi - \lambda_1) \ln \frac{1}{\varepsilon} + O(1).$$

By symmetry,

$$I_{\lambda_1, \lambda_2}(-u_\varepsilon) = 2(8\pi - \lambda_2) \ln \frac{1}{\varepsilon} + O(1).$$

In particular, it follows that whenever $\max\{\lambda_1, \lambda_2\} > 8\pi$, the functional I_{λ_1, λ_2} is unbounded below. Consequently, for all $(\lambda_1, \lambda_2) \in \Lambda$, I_{λ_1, λ_2} has a mountain pass structure. $\square$

Remark. The unboundedness of I_{λ_1, λ_2} when $\max\{\lambda_1, \lambda_2\} > 8\pi$ was derived by Ohtsuka and Suzuki [6] from certain properties of a functional related to the Moser-Trudinger inequality (6). However, we have proved it directly, since we shall use some properties of the function u_ε in the next section.

3 Proof of Theorem 1

Using Struwe's "monotonicity trick" [9] together with the blow-up analysis for (1) developed by Ohtsuka and Suzuki [6], we first prove the following result.

Proposition 2. *For a.e. $(\lambda_1, \lambda_2) \in \Lambda$ there exists a nonzero critical point for I_{λ_1, λ_2} .*

In order to prove Proposition 2, we begin by setting a minimax argument. In view of Proposition 1 and Lemma 3, there exists $\bar{u} = \bar{u}(\lambda_1, \lambda_2) \in E$ satisfying $\|\bar{u}\| \geq 1$ and $I_{\lambda_1, \lambda_2}(\bar{u}) < 0$. We consider the set of paths

$$\Gamma = \{\gamma \in C([0, 1], E) : \gamma(0) = 0, \gamma(1) = \bar{u}\}.$$

Then, the value

$$c(\lambda_1, \lambda_2) = \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} I_{\lambda_1, \lambda_2}(\gamma(t))$$

is finite. More precisely, the following estimate holds.

Lemma 4. *For every $\omega > 0$ there exists $\rho_\omega > 0$ independent of (λ_1, λ_2) such that*

$$c(\lambda_1, \lambda_2) \geq \frac{\rho_\omega^2}{2} \left(1 - \frac{\lambda_1 + \lambda_2}{\mu_1(M)|M|} - \omega \right).$$

Proof. By Taylor expansion of G at 0, where G is the functional defined in (3), in view of (4) we have:

$$G(u) = \frac{\int_M u^2 dv_g}{|M|} + o(\|u\|^2).$$

Hence, recalling that $\lambda_1 + \lambda_2 < \mu_1(M)|M|$ and the inequality (7), we may write

$$\begin{aligned} I_{\lambda_1, \lambda_2}(u) &= \frac{1}{2} \int_M |\nabla_g u|^2 dv_g - \frac{\lambda_1 + \lambda_2}{2|M|} \int_M u^2 dv_g + (\lambda_1 + \lambda_2) \circ (\|u\|^2) \\ &\geq \frac{1}{2} \left(1 - \frac{\lambda_1 + \lambda_2}{\mu_1(M)|M|} \right) \|u\|^2 - \mu_1(M)|M| \circ (\|u\|^2). \end{aligned}$$

Hence, for every $\omega > 0$ there exists $\rho_\omega \in (0, 1)$ independent of (λ_1, λ_2) such that

$$I_{\lambda_1, \lambda_2}(u) \geq \frac{1}{2} \left(1 - \frac{\lambda_1 + \lambda_2}{\mu_1(M)|M|} - \omega \right) \|u\|^2$$

for every $\|u\| \leq \rho_\omega$, and in particular

$$I_{\lambda_1, \lambda_2}(u) \geq \frac{\rho_\omega^2}{2} \left(1 - \frac{\lambda_1 + \lambda_2}{\mu_1(M)|M|} - \omega \right)$$

for all $u \in E$ satisfying $\|u\| = \rho_\omega$. Since for every $\gamma \in \Gamma$ there exists $\bar{t}$ for which $\|\gamma(\bar{t})\| = \rho_\omega$, we have

$$\max_{t \in [0, 1]} I_{\lambda_1, \lambda_2}(\gamma(t)) \geq I_{\lambda_1, \lambda_2}(\gamma(\bar{t})) \geq \frac{\rho_\omega^2}{2} \left(1 - \frac{\lambda_1 + \lambda_2}{\mu_1(M)|M|} - \omega \right).$$

Now the claim follows recalling the definition of $c(\lambda_1, \lambda_2)$. $\square$

We note that for every fixed $(\lambda_1, \lambda_2) \in \Lambda \setminus \partial\Lambda$ the function $\lambda \mapsto c(\lambda_1 + \lambda, \lambda_2 + \lambda)$ is well-defined and monotone decreasing for all λ near 0. Therefore it is differentiable at a.e. λ . Consequently, the function $\varepsilon \mapsto c(\lambda_1 + \varepsilon, \lambda_2 + \varepsilon)$ is differentiable at $\varepsilon = 0$ for a.e. $(\lambda_1, \lambda_2) \in \Lambda \setminus \partial\Lambda$. In what follows, such a $(\lambda_1, \lambda_2) \in \Lambda$ will be fixed, and to simplify notation, we set:

$$\begin{aligned} I &= I_{\lambda_1, \lambda_2}, \\ I_\varepsilon &= I_{\lambda_1 + \varepsilon, \lambda_2 + \varepsilon} = I - \varepsilon(G(u) + G(-u)) \\ c &= c_{\lambda_1, \lambda_2} \\ c_\varepsilon &= c(\lambda_1 + \varepsilon, \lambda_2 + \varepsilon) \\ c' &= \left. \frac{dc_\varepsilon}{d\varepsilon} \right|_{\varepsilon=0}, \end{aligned}$$

where the functional G is defined (3). We consider the set $X_\varepsilon = X_\varepsilon(\lambda_1, \lambda_2)$ defined by

$$X_\varepsilon = \left\{ u \in E : \begin{array}{l} u = \gamma(t) \text{ for some } \gamma \in \Gamma \text{ s.t. } \max_{t \in [0, 1]} I(\gamma(t)) \leq c + \varepsilon \\ \text{and } I_\varepsilon(u) \geq c_\varepsilon - \varepsilon \end{array} \right\}.$$

The following estimate holds:

Lemma 5. *Let $(\lambda_1, \lambda_2) \in \Lambda$ be such that $c'(\lambda_1, \lambda_2)$ exists. Then there exist constants $\eta > 0$, $C_1 > 0$ such that, for every $\varepsilon \in (0, \eta)$ and for every $u \in X_\varepsilon$, there holds:*

$$(i) \quad \|u\| \leq C_1$$

$$(ii) \quad c - (|c'| + 2)\varepsilon \leq I(u) \leq c + \varepsilon.$$

Proof. Since c' exists, we have

$$c_\varepsilon = c + c'\varepsilon + o(\varepsilon).$$

Proof of (i). For every $u \in X_\varepsilon$ we have, as $\varepsilon \rightarrow 0$:

$$\begin{aligned} 0 \leq G(u) + G(-u) &= \frac{I(u) - I_\varepsilon(u)}{\varepsilon} \leq \frac{c + \varepsilon - (c_\varepsilon - \varepsilon)}{\varepsilon} \\ &= -c' + 2 + o(1) = |c'| + 2 + o(1) \end{aligned}$$

and therefore there exists $\eta > 0$ such that

$$G(u) + G(-u) \leq |c'| + 3 \quad \forall u \in X_\varepsilon, \quad \forall \varepsilon \in (0, \eta).$$

It follows that

$$\begin{aligned} \|u\|^2 &= 2[I(u) + \lambda_1 G(u) + \lambda_2 G(-u)] \leq 2[c + \varepsilon + \max\{\lambda_1, \lambda_2\}(G(u) + G(-u))] \\ &\leq 2[c + \eta + \mu_1(M)|M|(|c'| + 3)] \end{aligned}$$

and therefore the asserted estimate (i) holds with $C_1^2 = 2[c + \eta + \mu_1(M)|M|(|c'| + 3)]$.

Proof of (ii). We have, recalling the monotonicity property of I_ε :

$$I(u) \geq I_\varepsilon(u) \geq c_\varepsilon - \varepsilon = c + c'\varepsilon + o(\varepsilon) - \varepsilon \geq c - (|c'| + 2)\varepsilon.$$

Since $I(u) \leq c + \varepsilon$ by definition of X_ε , (ii) is also established. $\square$

Now we show that if $c'(\lambda_1, \lambda_2)$ exists, then X_ε necessarily contains a bounded Palais-Smale sequence.

Lemma 6. *Let $(\lambda_1, \lambda_2) \in \Lambda$ be such that $c'(\lambda_1, \lambda_2)$ exists. Then there exists a sequence $(u_n)_{n \in \mathbb{N}}$, $u_n \in E$, such that $\|u_n\| \leq C_1$, $I(u_n) \rightarrow c$, $\|I'(u_n)\| \rightarrow 0$.*

Proof. If not, there exists $\delta > 0$ such that for every $u \in E$ satisfying $\|u\| \leq C_1$ and $|I(u) - c| < \delta$ we have $\|I'(u)\| \geq \delta$. In order to derive a contradiction, we need a suitable deformation. Let $\varepsilon_n \rightarrow 0$ and let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth cutoff function satisfying $0 \leq \varphi \leq 1$, $\varphi(s) = 1$ for all $s \geq -1$, $\varphi(s) = 0$ for all $s \leq -2$, and let

$$\Phi_n(u) = \varphi\left(\frac{I_{\varepsilon_n}(u) - c_{\varepsilon_n}}{\varepsilon_n}\right).$$

Then $\Phi_n(u) = 1$ for all u such that $I_{\varepsilon_n}(u) \geq c_{\varepsilon_n} - \varepsilon_n$ and in view of Lemma 4 we have $\Phi_n(0) = \Phi_n(\bar{u}) = 0$ for sufficiently large n . Consequently, setting

$$\tilde{\gamma}(t) = \gamma(t) - \sqrt{\varepsilon_n} \Phi_n(\gamma(t)) \frac{I'_{\varepsilon_n}(\gamma(t))}{\|I'_{\varepsilon_n}(\gamma(t))\|},$$

we have $\tilde{\gamma} \in \Gamma$ for every $\gamma \in \Gamma$. For every $n \in \mathbb{N}$, let $\gamma_n \in \Gamma$ be such that $\max_{t \in [0,1]} I(\gamma_n(t)) \leq c + \varepsilon_n$. We note that in view of Lemma 5, for all $u \in X_{\varepsilon_n}$ with sufficiently large n we have $\|u\| \leq C_1$, $|I(u) - c| < \delta$ and therefore, by assumption, we have $\|I'(u)\| \geq \delta$.

CLAIM. For sufficiently large n , we have

$$(13) \quad \max_{t \in [0,1]} I_{\varepsilon_n}(\tilde{\gamma}_n(t)) = \max_{\gamma_n(t) \in X_{\varepsilon_n}} I_{\varepsilon_n}(\tilde{\gamma}_n(t)).$$

Proof of Claim. By definition of c_{ε_n} we have

$$(14) \quad \max_{t \in [0,1]} I_{\varepsilon_n}(\tilde{\gamma}_n(t)) = \max_{\{t: I_{\varepsilon_n}(\tilde{\gamma}_n(t)) \geq c_{\varepsilon_n} - \varepsilon_n/2\}} I_{\varepsilon_n}(\tilde{\gamma}_n(t)).$$

We note that in view of (5) we have, for every $u, v \in E$:

$$\begin{aligned} \langle I''_{\varepsilon_n}(u)v, v \rangle &= \int_M |\nabla_g v|^2 dv_g - (\lambda_1 + \varepsilon_n) \langle G''(u)v, v \rangle - (\lambda_2 + \varepsilon_n) \langle G''(-u)v, v \rangle \\ &\leq \int_M |\nabla_g v|^2 dv_g \end{aligned}$$

and therefore, an expansion to the second order of I_{ε_n} yields

$$\begin{aligned} I_{\varepsilon_n}(u+v) &= I_{\varepsilon_n}(u) + I'_{\varepsilon_n}(u)v + \frac{1}{2} \langle I''_{\varepsilon_n}(u+sv)v, v \rangle \\ (15) \quad &\leq I_{\varepsilon_n}(u) + I'_{\varepsilon_n}(u)v + \frac{1}{2} \|v\|^2, \end{aligned}$$

where $s \in (0,1)$. Using (15), for all $t \in [0,1]$ such that $I_{\varepsilon_n}(\tilde{\gamma}_n(t)) > c_{\varepsilon_n} - \varepsilon_n/2$ we estimate:

$$\begin{aligned} c_{\varepsilon_n} - \frac{\varepsilon_n}{2} &< I_{\varepsilon_n}(\tilde{\gamma}_n(t)) \\ &\leq I_{\varepsilon_n}(\gamma_n(t)) + I'_{\varepsilon_n}(\gamma_n(t))(\tilde{\gamma}_n(t) - \gamma_n(t)) + \frac{1}{2} \|\tilde{\gamma}_n(t) - \gamma_n(t)\|^2 \\ &= I_{\varepsilon_n}(\gamma_n(t)) - \sqrt{\varepsilon_n} \Phi_n(\gamma_n(t)) \|I'_{\varepsilon_n}(\gamma_n(t))\| + \frac{\varepsilon_n}{2} \Phi_n^2(\gamma_n(t)) \\ &\leq I_{\varepsilon_n}(\gamma_n(t)) + \frac{\varepsilon_n}{2}. \end{aligned}$$

It follows that

$$I_{\varepsilon_n}(\gamma_n(t)) \geq c_{\varepsilon_n} - \varepsilon_n,$$

that is, $\gamma_n(t) \in X_{\varepsilon_n}$. We have obtained that

$$\left\{ t : I_{\varepsilon_n}(\tilde{\gamma}_n(t)) > c_{\varepsilon_n} - \frac{\varepsilon_n}{2} \right\} \subset \{ t : \gamma_n(t) \in X_{\varepsilon_n} \}.$$

Now, in view of the above and of (14), the asserted equivalence (13) follows and the claim is established.

Now we recall that for every t such that $\gamma_n(t) \in X_{\varepsilon_n}$, we have $\|\gamma_n(t)\| \leq C_1$, where C_1 is the constant obtained in Lemma 5. Consequently, we obtain from Lemma 1 that for every such t the following estimate holds:

$$(16) \quad \|I'_{\varepsilon_n}(\gamma_n(t))\| \geq \|I'(\gamma_n(t))\| - \varepsilon_n \|G'(\gamma_n(t))\| \geq \delta - \varepsilon_n C_M \exp \left\{ \frac{1}{8\pi} C_1^2 \right\} \geq \frac{\delta}{2},$$

for sufficiently large n . Moreover, for every t such that $\gamma_n(t) \in X_{\varepsilon_n}$ we also have $\Phi_n(\gamma_n(t)) = 1$ and $I(\gamma_n(t)) \leq c + \varepsilon_n$. Therefore, using (15) and (16) and the monotonicity property $I_{\varepsilon_n}(\gamma_n(t)) \leq I(\gamma_n(t))$, we estimate for all $\gamma_n(t) \in X_{\varepsilon_n}$:

$$\begin{aligned}
I_{\varepsilon_n}(\tilde{\gamma}_n(t)) &\leq I_{\varepsilon_n}(\gamma_n(t)) - \sqrt{\varepsilon_n} \Phi_n(\gamma_n(t)) \|I'_{\varepsilon_n}(\gamma_n(t))\| + \frac{\varepsilon_n}{2} \Phi_n(\gamma_n(t))^2 \\
&= I_{\varepsilon_n}(\gamma_n(t)) - \sqrt{\varepsilon_n} \|I'_{\varepsilon_n}(\gamma_n(t))\| + \frac{\varepsilon_n}{2} \\
&\leq I(\gamma_n(t)) - \frac{\delta \sqrt{\varepsilon_n}}{2} + \frac{\varepsilon_n}{2} \leq c + \varepsilon_n - \frac{\delta \sqrt{\varepsilon_n}}{2} + \frac{\varepsilon_n}{2} \\
&= c_{\varepsilon_n} - \frac{\delta \sqrt{\varepsilon_n}}{2} + \left(|c'| + \frac{3}{2} + o(1) \right) \varepsilon_n \\
&\leq c_{\varepsilon_n} - \frac{\delta \sqrt{\varepsilon_n}}{4}.
\end{aligned}$$

Therefore, in view of (13), we derive

$$\max_{t \in [0,1]} I_{\varepsilon_n}(\tilde{\gamma}_n(t)) = \max_{\gamma_n(t) \in X_{\varepsilon_n}} I_{\varepsilon_n}(\tilde{\gamma}_n(t)) \leq c_{\varepsilon_n} - \frac{\delta \sqrt{\varepsilon_n}}{4}$$

which in turn implies that

$$c_{\varepsilon_n} \leq c_{\varepsilon_n} - \frac{\delta \sqrt{\varepsilon_n}}{4},$$

a contradiction. It follows that such a δ does not exist, and that there necessarily exists a bounded Palais-Smale sequence at the level $c = c(\lambda_1, \lambda_2)$. $\square$

Proof of Proposition 2. Fix (λ_1, λ_2) so that $c(\lambda_1 + \lambda, \lambda_2 + \lambda)$ is differentiable with respect to λ at $\lambda = 0$. Let (u_n) be the bounded Palais-Smale sequence for $I = I_{\lambda_1, \lambda_2}$ as obtained in Lemma 6. We claim that (u_n) converges strongly in H^1 to a solution $u \not\equiv 0$ for problem (1). Indeed, since $\|u_n\| \leq C_1$, there exists $u \in E$ such that $u_n \rightharpoonup u$ weakly in H^1 , strongly in L^p for every $p \in [1, +\infty)$ and a.e. in M . Furthermore, $e^{u_n} \rightarrow e^u$ strongly in L^p for every $p \in [1, +\infty)$ and $e^{-u_n} \rightarrow e^{-u}$ strongly in L^p for every $p \in [1, +\infty)$. Therefore, taking limits in the equation

$$\int_M \nabla_g u_n \cdot \nabla_g \phi \, dv_g = \lambda_1 \frac{\int_M e^{u_n} \phi \, dv_g}{\int_M e^{u_n} \, dv_g} - \lambda_2 \frac{\int_M e^{-u_n} \phi \, dv_g}{\int_M e^{-u_n} \, dv_g} + o(1) \quad \forall \phi \in E,$$

we see that u is a solution for (1). Moreover, since $\|I'(u_n)\| \rightarrow 0$,

$$o(1)(\|u_n - u\|) = \langle I'(u_n), u_n - u \rangle = \|u_n - u\|^2 + o(1)$$

and therefore, $u_n \rightarrow u$ strongly in H^1 . By continuity and in view of Lemma 4 we derive, as $n \rightarrow +\infty$

$$I(u_n) \rightarrow I(u) = c > 0$$

and consequently $u \not\equiv 0$. $\square$

Let $(\lambda_1^*, \lambda_2^*) \in \Lambda$ and let $(\lambda_{1,n}, \lambda_{2,n}) \in \Lambda$, $(\lambda_{1,n}, \lambda_{2,n}) \rightarrow (\lambda_1^*, \lambda_2^*)$ be such that problem (1) admits corresponding nonzero solutions (u_n) as obtained in Proposition 2. We note that, in view of Lemma 4 with

$$\omega = \bar{\omega} = \frac{1}{2} \left(1 - \frac{\lambda_1^* + \lambda_2^*}{\mu_1(M)|M|} \right)$$

we have, as $n \rightarrow +\infty$,

$$(17) \quad c(\lambda_{1,n}, \lambda_{2,n}) \geq \frac{\rho_\omega^2}{4} \left(1 - \frac{\lambda_1^* + \lambda_2^*}{\mu_1(M)|M|} \right) + o(1).$$

Following the notation in [6], we denote by $\mathcal{S}_1, \mathcal{S}_2$ the blow-up sets of (u_n) and $(-u_n)$, respectively. Namely, we set

$$\begin{aligned} \mathcal{S}_1 &= \{x \in M : \exists x_n \rightarrow x \text{ s.t. } u_n(x_n) \rightarrow +\infty\} \\ \mathcal{S}_2 &= \{x \in M : \exists x_n \rightarrow x \text{ s.t. } u_n(x_n) \rightarrow -\infty\}. \end{aligned}$$

Moreover, we set

$$\mu_{1,n} = \lambda_{1,n} \frac{e^{u_n}}{\int_M e^{u_n} dv_g}, \quad \mu_{2,n} = \lambda_{2,n} \frac{e^{-u_n}}{\int_M e^{-u_n} dv_g}.$$

Since $\int_M d\mu_{i,n} = \lambda_{i,n}$ for $i = 1, 2$, there exist Radon measures μ_i , $i = 1, 2$, such that $\mu_{i,n} \rightharpoonup \mu_i$, $i = 1, 2$, weakly in the sense of measures. In particular,

$$(18) \quad \lambda_{i,n} = \int_M d\mu_{i,n} \rightarrow \lambda_i^* = \int_M d\mu_i \quad i = 1, 2.$$

At this point the following blow-up analysis from [6] is a key step.

Proposition 3 ([6]). *For the solution sequence (u_n) exactly one of the following alternatives occurs:*

- (i) *compactness: $\mathcal{S}_1 \cup \mathcal{S}_2 = \emptyset$. Up to subsequences, $u_n \rightarrow u^* \in E$, where u^* is solution to (1) with $\lambda_1 = \lambda_1^*$ and $\lambda_2 = \lambda_2^*$.*
- (ii) *one-sided concentration: $\mathcal{S}_i \neq \emptyset$ and $\mathcal{S}_j = \emptyset$, $j \neq i$. In this case $\mu_i = \sum_{x_0 \in \mathcal{S}_i} 8\pi \delta_{x_0}$. In particular, $\lambda_i^* = 8\pi \text{card}(\mathcal{S}_i)$.*
- (iii) *two-sided concentration: $\mathcal{S}_1 \neq \emptyset$ and $\mathcal{S}_2 \neq \emptyset$. In this case, $\mu_i = r_i + \sum_{x_0 \in \mathcal{S}_i} m_i(x_0) \delta_{x_0}$ with $r_i \geq 0$, $r_i \in L^1(M) \cap L^\infty_{\text{loc}}(M \setminus \mathcal{S}_i)$ and $m_i(x_0) \geq 4\pi$. Furthermore,*
 - (iii-a) *If there exists $x_0 \in \mathcal{S}_i \setminus \mathcal{S}_j$, then $r_i \equiv 0$ and $m_i(x_0) = 8\pi$.*
 - (iii-b) *For every $x_0 \in \mathcal{S}_1 \cap \mathcal{S}_2$ the following relation holds: $(m_1(x_0) - m_2(x_0))^2 = 8\pi(m_1(x_0) + m_2(x_0))$.*

At this point we can complete the proof of Theorem 1.

Proof of Theorem 1. We begin by showing that $\text{card}(\mathcal{S}_1 \cap \mathcal{S}_2) = \emptyset$. Indeed, in view of (18) and Proposition 3-(iii), we obtain

$$\lambda_1^* + \lambda_2^* = \int_M d\mu_1 + \int_M d\mu_2 \geq m_1(x_0) + m_2(x_0).$$

On the other hand, the following property is elementary:

$$\begin{aligned} \min \{x + y : x, y \geq 4\pi, (x - y)^2 = 8\pi(x + y)\} &= \min \left\{ \eta : \eta \geq 8\pi + |\xi|, \eta = \frac{\xi^2}{8\pi} \right\} \\ &= 4(3 + \sqrt{5}). \end{aligned}$$

It follows from Proposition 3-(iii-b) that if $x_0 \in \mathcal{S}_1 \cap \mathcal{S}_2$, then

$$\lambda_1^* + \lambda_2^* \geq 4(3 + \sqrt{5})\pi > 16\pi > \mu_1(M)|M|$$

and therefore $(\lambda_1^*, \lambda_2^*) \notin \Lambda$, a contradiction. Now suppose that $\mathcal{S}_1 = \mathcal{S}_1 \setminus \mathcal{S}_2 \neq \emptyset$. Then, in view of Proposition 3-(iii-a) we obtain $r_1 \equiv 0$ and $\mu_1 = 8\pi \sum_{x_0 \in \mathcal{S}_1} \delta_{x_0}$ and thus (18) implies $\lambda_1^* = 8\pi \text{card}(\mathcal{S}_1) > 0$ and again we obtain $(\lambda_1^*, \lambda_2^*) \notin \Lambda$, a contradiction. Similarly, we see that $\mathcal{S}_2 = \emptyset$. Hence, $\mathcal{S}_1 \cup \mathcal{S}_2 = \emptyset$. Therefore, alternative (i) of Proposition 3 holds, and there exists a solution $u^* \in E$ to problem (1) such that $u_n \rightarrow u^*$ strongly in H^1 . By continuity, $I_{\lambda_1, n, \lambda_2, n}(u_n) \rightarrow I_{\lambda_1^*, \lambda_2^*}(u^*)$. In particular, in view of (17), we have

$$I_{\lambda_1^*, \lambda_2^*}(u^*) \geq \liminf_{n \rightarrow +\infty} c(\lambda_{1, n}, \lambda_{2, n}) \geq \frac{\rho_\omega^2}{4} \left(1 - \frac{\lambda_1^* + \lambda_2^*}{\mu_1(M)|M|} \right) > 0$$

and therefore $u^* \neq 0$. □

References

- [1] H. Brezis and F. Merle, Uniform estimates and blow-up behavior for solutions to $-\Delta u = V(x)e^u$ in two dimensions, *Comm. Partial Differential Equations* **16** (1991), 1223–1253.
- [2] D. Chae, H. Ohtsuka and T. Suzuki, Some existence results for solutions to $SU(3)$ Toda system, *Calc. Var.* **24** No. 4 (2005), 403–429.
- [3] L. Jeanjean, On the existence of a bounded Palais-Smale sequence and application to a Landesman-Lazer type problem, *Proc. Roy. Soc. Edinburgh* **129A** (1999), 787–809.
- [4] G. Joyce, D. Montgomery, Negative temperature states for the two-dimensional guiding-centre plasma, *J. Plasma Phys.* **10** (1973), 107–121.
- [5] M. Lucia and M. Nolasco, $SU(3)$ Chern-Simons Vortex Theory and Toda Systems, *J. Differential Equations* **184** (2002), 443–474.
- [6] H. Ohtsuka and T. Suzuki, Mean field equation for the equilibrium turbulence and a related functional inequality, *Adv. Differential Equations* **11** No. 3 (2006), 281–304.
- [7] Y.B. Pointin and T.S. Lundgren, Statistical mechanics of two-dimensional vortices in a bounded container, *Phys. Fluids* **19** (1976), 1459–1470.
- [8] K. Sawada, T. Suzuki and F. Takahashi, Mean field equation for equilibrium vortices with neutral orientation, preprint.
- [9] M. Struwe, The existence of surfaces of constant mean curvature with free boundaries, *Acta Math.* **160** (1988), 19–64.
- [10] M. Struwe and G. Tarantello, On multivortex solutions in Chern-Simons gauge theory, *Boll. Unione Mat. Ital. Sez. B* (8)**1** (1998), 109–121.